

hepth@xxx/9203039

LPTENS-92/07

ENSLAPP-L-370/92

March 1992

BLACK HOLES FROM NON-ABELIAN TODA THEORIES

Jean-Loup GERVAIS

Laboratoire de Physique Théorique de l'École Normale Supérieure¹,
24 rue Lhomond, 75231 Paris CÉDEX 05, France.

and

Mikhail V. SVELTIEV²

Laboratoire de Physique Théorique de l'ENS Lyon,
46 Allée d'Italie, 69364 Lyon CÉDEX 07, France.

Abstract

Non-abelian Toda theories are shown to provide exactly solvable conformal systems in the presence of a black hole. They correspond to gauged WZNW models where the gauge group is nilpotent, and are thus basically different from the ones currently considered, following Witten. The non-abelian Toda potential gives a cosmological term which may be integrated exactly at the classical level.

¹Unité Propre du Centre National de la Recherche Scientifique, associée à l'École Normale Supérieure et à l'Université de Paris-Sud.

² On leave of absence from the Institute for High Energy Physics, 142284, Protvino, Moscow region, Russia.

From the beginning the conformal field theories in two-dimensions (2D CFT) have been considered using two different viewpoints. The celebrated BPZ approach [1], made a systematic use of Ward identities assuming that renormalization gives simple numerical modifications of the coupling constants. It came out at a time where the authors of ref.[2] had already started to solve the Liouville theory. The latter work was the beginning of the modern investigations based on the connection between 2D CFT and integrable systems. In particular, the authors of papers [3] presented a classification of the CFT associated with the Toda systems, and of the related W-symmetries by putting them in correspondence with the simple finite-dimensional Lie algebras $\mathcal{G}$ over $\mathbf{C}$. In the present letter, we discuss the so-called non-abelian Toda theories. Most of these models are derivable from σ -model actions of the form (γ is the coupling constant)

$$\mathcal{S} = \frac{1}{\gamma} \int d_2z \sqrt{-h} \left\{ \frac{1}{2} G_{ij}(x) h^{\mu\nu} \partial_\mu x^i \partial_\nu x^j - T(x) + \Phi(x) R^{(2)} \right\}. \quad (1)$$

Greek indices are related to the world-sheet which has metric tensor $h^{\mu\nu}$, and scalar curvature $R^{(2)}$. Latin indices run from 1 to N . The functions $x^i(z^+, z^-)$ are bosonic fields which may be regarded as components of a string-position in a N -dimensional target space with metric $G_{ij}(x)$. From this viewpoint, T and Φ are the tachyon and dilaton fields respectively. In the present article, we shall mostly work with constant Minkowski 2D metric $h_{\mu\nu} = -\eta_{\mu\nu}$, in which case, the dilaton field only appears in the expression of the (improved) stress-energy tensor

$$\theta_{\mu\nu} \equiv 2 \frac{\delta \mathcal{S}}{\delta h^{\mu\nu}} \Big|_{h^{\mu\nu} = -\eta^{\mu\nu}} = T_{\mu\nu} + 2(\eta_{\mu\nu} \partial^2 - \partial_\mu \partial_\nu) \Phi, \quad (2)$$

where $T_{\mu\nu}$ is the canonical (Noether) stress-energy tensor with the flat 2D metric. As an example, the abelian Toda theories, considered in ref.[3] are associated with the Lagrangian

$$-\gamma \mathcal{L} = \frac{1}{2} \sum_{i,j=1}^N (w K^{(\mathcal{G})})_{ij} \frac{\partial x^i}{\partial z_+} \frac{\partial x^j}{\partial z_-} + \sum_{i=1}^N \prod_{j=1}^N w_i \exp \left(K_{ij}^{(\mathcal{G})} x^j \right), \quad (3)$$

where G_{ij} coincides with the symmetrized Cartan matrix of a simple Lie algebra $\mathcal{G}$ of rank N . Explicitly, denoting by $\vec{\alpha}_i$ a set of primitive roots of $\mathcal{G}$, $K_{ij}^{(\mathcal{G})} = (\vec{\alpha}_i \cdot \vec{\alpha}_j) 2/\alpha_j^2$ is the Cartan matrix, and $w \equiv \text{diag}(w_1, \dots, w_N)$, $w_i = 2/\alpha_i^2$. These theories are of the type Eq.1 with

$$T \equiv \sum_{i=1}^N \prod_{j=1}^N w_i \exp \left(K_{ij}^{(\mathcal{G})} x^j \right), \quad \Phi = \sum_{i=1}^N \omega_i x^i. \quad (4)$$

Clearly, the coefficients ω_i in T may be changed to any value by shift of the fields x . These defines N cosmological constants which may be re-introduced whenever needed. The expression for Φ follows from the study[3] of the conformal invariance properties of these systems. The fields x^i satisfy the equation

$$\partial^2 x^i / \partial z_+ \partial z_- = \prod_j \exp \left(K_{ij}^{(\mathcal{G})} x^j \right), \quad 1 \leq i \leq N; \quad (5)$$

It is natural to characterize such systems as being abelian since, in the framework of an algebraic approach, (see e.g. review [4]), they are associated with the canonical

gradation $\mathcal{G} = \oplus_{m \in \mathbf{Z}} \mathcal{G}_m$ of the simple Lie algebras, for which the subalgebra $\mathcal{G}_0$ is abelian. Indeed, the canonical gradation is such that the step operators associated with positive (resp. negative) simple roots have grading 1 (resp. -1), and $\mathcal{G}_0$ coincides with the Cartan subalgebra. In other words, this canonical gradation is related with the principal $sl(2, \mathbf{C})$ subalgebra of $\mathcal{G}$.

The above example is somewhat trivial from the target-space viewpoint since the metric G^{ij} is constant. The point of the present letter is to remove this restriction, by considering the so-called non-abelian versions of the Toda systems. These theories, constructed in [5], correspond to non-canonical gradations of simple Lie algebras where the subalgebra $\mathcal{G}_0$ is non-abelian. Such $\mathbf{Z}$ -gradations of $\mathcal{G}$ are related with the non-principal embeddings of $sl(2, \mathbf{C})$ in $\mathcal{G}$. We shall put forward the examples of the Lie algebras B_{N-1} , since they give integrable theories with black-hole background metrics similar to the one recently popularized by Witten[6][7] [8] [9] [10]. Associated with a maximal subalgebra D_{N-1} (of maximal rank) in the Lie algebra B_{N-1} , we have the Lagrangian

$$\begin{aligned}
-\gamma \mathcal{L} = & \frac{1}{2} \left[\sum_{j,k=1}^{N-1} K_{jk}^{(D_{N-1})} \frac{\partial x^j}{\partial z_+} \frac{\partial x^k}{\partial z_-} - \tanh^2 \left(\frac{x^{N-1} - x^{N-2}}{2} \right) \frac{\partial x^N}{\partial z_+} \frac{\partial x^N}{\partial z_-} \right] \\
& + \sum_{i=1}^{N-1} \prod_{j=1}^{N-1} \exp \left(K_{ij}^{(D_{N-1})} x^j \right). \quad (6)
\end{aligned}$$

Note that there is no useful non-abelian version of the Toda lattice for the series A_N . In this case, the zero-grading part is given by $\mathcal{G}_0 = gl(1) \oplus \cdots \oplus gl(1) \oplus B_1$, where the one-dimensional linear algebra $gl(1)$ appears $N-1$ times. The x -dependent part of the metric involves the hyperbolic-tangent-square function which is familiar in the 2D black-hole game. In particular if we choose $N=3$, we find

$$-\gamma \mathcal{L} = \frac{1}{2} \left[2 \sum_{i=1}^2 \frac{\partial x^i}{\partial z_+} \frac{\partial x^i}{\partial z_-} - \tanh^2 \left(\frac{x^1 - x^2}{2} \right) \frac{\partial x^3}{\partial z_+} \frac{\partial x^3}{\partial z_-} \right] + \sum_{i=1}^2 \exp(2x^i) \quad (7)$$

In order to clarify the black-hole aspect, it is convenient to change field variables. Let

$$\phi = (x^1 + x^2)/2, \quad r = (x^1 - x^2)/2, \quad \theta = x^3/2. \quad (8)$$

One gets

$$-\frac{\gamma}{2} \mathcal{L} = \frac{\partial \phi}{\partial z_+} \frac{\partial \phi}{\partial z_-} + \frac{\partial r}{\partial z_+} \frac{\partial r}{\partial z_-} - \tanh^2(r) \frac{\partial \theta}{\partial z_+} \frac{\partial \theta}{\partial z_-} + \cosh(2r) e^{2\phi}. \quad (9)$$

The target-space metric is

$$G = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -\tanh^2 r \end{pmatrix} \quad (10)$$

one sees that the first component, which corresponds to the ϕ variable, defines a subspace that decouples from the rest from the viewpoint of Riemannian geometry. In the (r, θ) space we have $G = \begin{pmatrix} 1 & 0 \\ 0 & -\tanh^2 r \end{pmatrix}$, which coincides with Witten's black-hole

metric[10] exactly. As is well known by now the corresponding manifold is a cigar with radial variable r , and angle θ . Letting

$$u = \sinh(r)e^{-\theta}, \quad v = -\sinh(r)e^{\theta}, \quad (11)$$

one finds the standard form $-dudv/(1-uv) = dr^2 - d\theta^2 \tanh^2 r$. The action becomes

$$\frac{\gamma}{2}\mathcal{L} = \frac{1}{(1-uv)}(\partial_+ u \partial_- v + \partial_- u \partial_+ v) - \partial_+ \phi \partial_- \phi - e^{2\phi}(1-2uv). \quad (12)$$

Note that there exists another non-canonical gradation of B_N , with $\mathcal{G}_0 = gl(1) \oplus B_{N-1}$, for which the corresponding non-abelian version of the Toda systems possess the Lagrangian

$$\frac{\gamma}{2}\mathcal{L} = \frac{1}{(1-\vec{u} \cdot \vec{v})}(\partial_+ \vec{u} \cdot \partial_- \vec{v} + \partial_- \vec{u} \cdot \partial_+ \vec{v}) - \partial_+ \phi \partial_- \phi - e^{2\phi}(1-2\vec{u} \cdot \vec{v}), \quad (13)$$

where $\vec{u}$ and $\vec{v}$ are $(N-1)$ -dimensional vectors. It represents a black higher-dimensional manifold.

For completeness, let us next recall some standard results on the systems we are discussing. The basic point of the Toda theories — abelian as well as non-abelian — is that their dynamical equations, no matter how complicated, are derivable from flatness conditions

$$[\partial_+ - \mathcal{A}_+, \partial_- - \mathcal{A}_-] \equiv \partial_- \mathcal{A}_+ - \partial_+ \mathcal{A}_- + [\mathcal{A}_+, \mathcal{A}_-] = 0, \quad (14)$$

which allow to obtain the general solutions in closed form[4][5]. The Lax pair $\mathcal{A}_{\pm}$ is systematically constructed once the Lie algebra, the gradation, and the grading spectrum of $\mathcal{A}_{\pm}$ are chosen[4][5]. These results were re-derived recently[11] by proving that the Toda theories may be obtained by conformal reduction of WZNW models. Let us recall some basic point about Toda solutions, following refs.[4][5], but stressing the WZNW connection of ref.[11]. Consider a finite-dimensional Lie group $\mathbf{G}$, with Lie algebra $\mathcal{G}$, and a gradation $\mathcal{G} = \oplus_{m \in \mathbf{Z}} \mathcal{G}_m$, with $[\mathcal{G}_m, \mathcal{G}_n] \in \mathcal{G}_{m+n}$. There is a grading operator H , such that $[H, \mathcal{G}_m] = 2m\mathcal{G}_m$. Associated with this grading, we have the (so-called modified) Gauss decomposition of the Lie group $\mathbf{G}$, namely, any regular element ω may be written as

$$\omega \equiv \omega_+ \omega_0 \omega_-, \quad \text{with } \omega_{\epsilon} \in \mathbf{G}_{\epsilon}, \quad \epsilon = \pm, 0. \quad (15)$$

$\mathbf{G}_0$, $\mathbf{G}_{\pm}$ are the subgroups generated by $\mathcal{G}_0$, and $\mathcal{G}_{\pm} \equiv \oplus_{m>0} \mathcal{G}_{\pm m}$, respectively. With the simplest choice of gradation spectrum, the Lax pair is given by³

$$\begin{aligned} \mathcal{A}_- &\equiv \partial_- g_0^{-1} g_0 - M_-, \\ \mathcal{A}_+ &\equiv -g_0^{-1} M_+ g_0, \end{aligned} \quad (16)$$

where $g_0 \in \mathbf{G}_0$, and where $M_{\pm}$ are, respectively, fixed elements of $\mathcal{G}_{\pm 1}$, such that the action of $\mathcal{G}_0$ on them generates the whole $\mathcal{G}_{\pm 1}$. The connection with the WZNW model goes as follows[11]. Let $\omega_0 = g_0^{-1}$, and define $\omega_{\pm}$ as solutions of the equations

$$\omega_+^{-1} \partial_+ \omega_+ = \omega_0 M_+ \omega_0^{-1}, \quad \omega_- \partial_- \omega_-^{-1} = \omega_0^{-1} M_- \omega_0. \quad (17)$$

³By convention, the differential operators $\partial_{\pm}$ only act on the first function on the right, unless parenthesis indicate otherwise.

Clearly, the solutions are such that $\omega_{\pm} \in \mathbf{G}_{\pm}$, and if we define $\omega \equiv \omega_+ \omega_0 \omega_-$, we have a Gauss decomposition of the type Eq.15. A straightforward calculation then shows[11] that the currents

$$\partial_- \omega \omega^{-1} = J, \quad \omega^{-1} \partial_+ \omega = -\bar{J}. \quad (18)$$

satisfy the equations

$$\partial_+ J = \partial_- \bar{J} = 0, \quad (19)$$

and it follows that ω is a solution of the WZNW equations. Moreover, one may verify that the currents are such that their projections $\Pi_{\pm}$ into $\mathcal{G}_{\pm}$ are given by

$$\Pi_-(J) = -M_-, \quad \Pi_+(\bar{J}) = -M_+. \quad (20)$$

Next, the general solution of the dynamical equation associated with Eq.16, that is

$$\partial_+ (\partial_- g_0^{-1} g_0) + [M_-, g_0^{-1} M_+ g_0] = 0 \quad (21)$$

has been known for quite some time[4][5][14]. It was recently connected[11] with the WZNW solution as follows. The general solution of Eq.19 is $\omega = \omega_L(z_-) \omega_R(z_+)$ with $J = \partial_- \omega_L \omega_L^{-1}$, $\bar{J} = -\omega_R^{-1} \partial_+ \omega_R^{-1}$. The group element g relevant for the Toda theory is $g \equiv \omega^{-1}$, and we have $g = g_L(z_+) g_R(z_-)$, with $g_L = \omega_R^{-1}$, $g_R = \omega_L^{-1}$ and $g = g_- g_0 g_+$, $g_{\pm} \equiv \omega_{\pm}^{-1}$. It next follows from Eqs.17 that, if we introduce the decompositions $g_L = g_{-L} g_{0L} g_{+L}$, $g_R = g_{-R} g_{0R} g_{+R}$, we have

$$\begin{aligned} \partial_+ g_{+L} &= -\bar{f}(z_+) g_{+L}, & \bar{f}(z_+) &\equiv g_{0L}^{-1} M_+ g_{0L} \in \mathcal{G}_+ \\ \partial_- g_{-R} &= g_{-R} f(z_-), & f(z_-) &\equiv g_{0R}^{-1} M_- g_{0R} \in \mathcal{G}_-, \end{aligned} \quad (22)$$

with solutions

$$\begin{aligned} g_{+L}(z) &= \sum_n (-1)^n \int^z dx_1 \cdots \int^{x_{n-1}} dx_n \bar{f}(x_1) \cdots \bar{f}(x_n), \\ g_{-R}(z) &= \sum_n \int^z dx_1 \cdots \int^{x_{n-1}} dx_n f(x_n) \cdots f(x_1), \end{aligned} \quad (23)$$

where only a finite number of terms are non-vanishing, since the Lie algebra considered is finite-dimensional. The solution of Eq.21 is obtained by computing the matrix-elements of g between highest-weight vectors $|\lambda\rangle$ which are annihilated by $\mathcal{G}_m$ for positive m , so that $\langle \lambda | g | \lambda \rangle = \langle \lambda | g_0 | \lambda \rangle$. One finds

$$\langle \lambda | g_0 | \lambda \rangle = \langle \lambda | g_{L0} g_{L+} g_{R-} g_{R0} | \lambda \rangle, \quad (24)$$

which, according to Eq.23, coincides with the general solution of refs.[4][5][14]. In practice, f and $\bar{f}$ are arbitrary elements of $\mathcal{G}_-$ and $\mathcal{G}_+$, respectively, functions of a single variable. In the same way as for the abelian case[3], they are classical analogues of the quantum screening operators. In general, the zero-grading part $\mathcal{G}_0$ involves a subalgebra $\mathcal{G}_0^0$, that commutes with $M_{\pm}$. It follows from Eq.21 that in g_0 , only the part that do not belong to $\mathcal{G}_0^0$ has a non-trivial dynamics. This terminates the short summary of general properties.

Returning to the case of the non-abelian B_2 Toda theory, it is illustrative to make use of the following explicit realization[15] of the Lie algebra B_2 . Consider two orthonormal vectors $\vec{e}_{1,2}$ in an Euclidian 2D vector space. The positive roots of B_2 are

$$\vec{\pi}_1 = \vec{e}_1 - \vec{e}_2, \quad \vec{\pi}_2 = \vec{e}_2, \quad \vec{\pi}_{12} = \vec{e}_1, \quad \vec{\pi}_{122} = \vec{e}_1 + \vec{e}_2, \quad (25)$$

where the first two are simple. Introduce five fermionic oscillators b_j , $j = \pm 1, \pm 2, 0$, with $[b_j, b_k^+] = \delta_{j,k}$. The B_2 generators are

$$\begin{aligned} h_1 &= b_1^+ b_1 - b_{-1}^+ b_{-1} - b_2^+ b_2 - b_{-2}^+ b_{-2}, \\ h_2 &= 2(b_2^+ b_2 - b_{-2}^+ b_{-2}) \\ E_{e_1 - e_2} &= b_1^+ b_2 - b_{-2}^+ b_{-1}, \\ E_{e_2} &= \sqrt{2}(b_2^+ b_0 - b_0^+ b_{-2}), \\ E_{e_1} &= \sqrt{2}(b_1^+ b_0 - b_0^+ b_{-1}), \\ E_{e_1 + e_2} &= b_1^+ b_{-2} - b_2^+ b_{-1}. \end{aligned} \quad (26)$$

The non-abelian Toda theory arises by choosing

$$H = 2(b_1^+ b_1 - b_{-1}^+ b_{-1}), \quad M_+ = E_{e_1}, \quad M_- = E_{e_1}^+. \quad (27)$$

These operators generate a non-canonical embedding of A_1 in B_2 . The gradation 0 part $\mathcal{G}_0$ is the linear span of $h_1, h_2, E_{e_2}, E_{e_2}^+$, and $\mathcal{G}_0^0$ is clearly generated by h_2 . The subspace $\mathcal{G}_1$ is the linear span of $E_{e_1 - e_2}, E_{e_1 + e_2}, E_{e_1}$. The first two are step operators corresponding to simple roots of D_2 . The parametrization of g_0 is

$$g_0 = \exp(a^+ E_{e_2}) \exp(a^- E_{e_2}^+) \exp(a_1 h_1 + a_2 h_2), \quad (28)$$

which gives

$$\begin{aligned} g_0^{-1} \partial_- g_0 &= \partial_- a_1 H / 2 + (\partial_- a_2 - \partial_- a_1 / 2 + \partial_- a^+ a^-) h_2 + e^{-2a_2 + a_1} \partial_- a^+ E_{e_2} + \\ &+ e^{2a_2 - a_1} [\partial_- a^- - \partial_- a^+ (a^-)^2] E_{e_2}^+. \end{aligned} \quad (29)$$

$$\begin{aligned} g_0^{-1} M_+ g_0 &= e^{-a_1} \left\{ -2a^+ e^{a_1 - 2a_2} E_{e_1 + e_2} + (1 + 2a^+ a^-) E_{e_1} + \right. \\ &\left. 2a^- (1 + a^+ a^-) e^{-a_1 + 2a_2} E_{e_1 - e_2} \right\}. \end{aligned} \quad (30)$$

The final equations, equivalent to Eq.21 are

$$\begin{aligned} \partial_+ \partial_- a_1 &= -2(1 + 2a^+ a^-) e^{-a_1}, \\ \partial_+ \partial_- a_2 + \partial_+ (\partial_- a^+ a^-) &= -(1 + 2a^+ a^-) e^{-a_1}, \\ \partial_+ \{ e^{a_1 - 2a_2} \partial_- a^+ \} &= 2a^+ e^{-2a_2}, \\ \partial_+ \{ e^{-a_1 + 2a_2} \partial_+ (-\partial_- a^+ (a^-)^2 + \partial_- a^-) \} &= 2a^- (1 + a^+ a^-) e^{-2a_1 + 2a_2}. \end{aligned} \quad (31)$$

These equations are equivalent to the field-equations derived from the action Eq.9 if we let

$$\phi = -a_1, \quad \sqrt{a^+ a^-} = \cosh(r/2), \quad (32)$$

$$\begin{aligned}
\partial_+ \theta / \sqrt{2} &= -(1 + a^+ a^-) \partial_+ \left\{ \ln \left(a^- e^{-a_1 + 2a_2} \right) \right\} + \partial_+ \left\{ \ln a^+ a^- \right\} / 2, \\
\partial_- \theta / \sqrt{2} &= \frac{1 + a^+ a^-}{1 + 2a^+ a^-} \partial_- \left\{ \ln \left(a^+ e^{a_1 - 2a_2} \right) \right\} - \partial_- \left\{ \ln a^+ a^- \right\} / 2.
\end{aligned} \tag{33}$$

Finally, one makes use of Eq.24 with the three possible highest-weight states

$$|\lambda_1\rangle = b_1^+ |0\rangle, \quad |\lambda_2\rangle = b_2^+ b_1^+ |0\rangle, \quad |\lambda_0\rangle = b_0^+ b_1^+ |0\rangle. \tag{34}$$

Combining the result with Eq.22 gives the exact solutions in terms of the screening fields f and $\bar{f}$ (see ref.[5] for details). Such is the classical Coulomb gas picture of the black-hole solution.

Our next topic is the connection with gauged WZNW models. Witten's solution is obtained from the $sl(2, R)$ – WZNW model by gauging the $gl(1, R)$ subgroup $\omega \rightarrow \exp(\alpha\sigma_3) \omega \exp(\alpha\sigma_3)$, α being an arbitrary function. It may be generalized straightforwardly by embedding this $sl(2, R)$ into larger groups. Its basic characteristic is that one gauges a diagonal one-parameter subgroup. In our explicit example, the non-abelian Toda theory was based on the Lie algebra B_{N-1} . Then the group of the corresponding WZNW model is the associated maximally-non-compact real form, that is, $so(N-1, N)$. As shown in ref.[11], the Toda theories (abelian or not) are obtained by gauging a nilpotent algebra instead of a diagonal one, and the gauge-group transformation is given by $\omega \rightarrow h_+ \omega h_-^{-1}$, where $h_{\pm} \in \mathbf{G}_{\pm}$, and $\mathbf{G}_{\pm}$ is specified by the gradation of the Lie algebra considered (see above). For instance, for $N = 2$ the action Eq.9 is obtained from the $so(1, 2)$ – WZNW model by gauging the transformation

$$\omega \rightarrow \exp(\alpha E_{e_1 - e_2} + \beta E_{e_1 + e_2} + \gamma E_{e_1}) \omega \exp(-\tilde{\alpha} E_{e_1 - e_2}^+ - \tilde{\beta} E_{e_1 + e_2}^+ - \tilde{\gamma} E_{e_1}^+), \tag{35}$$

where $\alpha, \dots, \tilde{\gamma}$ are arbitrary. Clearly this mechanism is more complicated than Witten's. However the dynamics is completely integrable, and we are able to obtain the solution in closed form to arbitrary order in the tachyon field $T \propto e^{2\phi}(1 - 2uv)$. Thus this particular tachyon field is exactly solved.

Now we come to the dilaton field Φ , and to the conformal invariance. At the purely classical level (following ref.[3]), Φ is determined from the requirement that $\theta_{\mu\nu}$ — which is defined by Eq.2 — be traceless, as a consequence of the equations of motions

$$\partial^2 x^i / \partial z_+ \partial z_- - \sum_j G^{ij}(x) \frac{\delta T(x)}{\delta x^j} + \sum_{j,l} \Gamma_{jl}^i(x) \frac{\partial x^j}{\partial z_+} \frac{\partial x^l}{\partial z_-} = 0. \tag{36}$$

Here $\Gamma_{jl}^i(x)$ are the Christoffel symbols associated with G . Conformal invariance requires that

$$\partial^2 \Phi(x) / \partial z_+ \partial z_- - T(x) = 0, \tag{37}$$

which is equivalent to

$$\sum_{i,j} \left(\partial_+ x^i \partial_- x^j \Phi_{;ij} + G^{ij} \frac{\delta \Phi}{\delta x^i} \frac{\delta T}{\delta x^j} \right) - T = 0$$

where semi-colons denote covariant derivatives with respect to x . For a given value of $z_{\pm}$, $\partial_{\pm} x^j$, and x^j are independant vectors. Since T and Φ are taken to be independant

of $\partial_{\pm}x^j$, it follows that we must have

$$\Phi_{;ij} = 0, \quad i, j = 1, \dots, N; \quad \sum_{\ell, m} G^{\ell m} \frac{\delta \Phi}{\delta x^{\ell}} \frac{\delta T}{\delta x^m} - T = 0. \quad (38)$$

The equations $v_{i;j} \equiv \partial v_i / \partial \phi^j - \Gamma_{ij}^l v_l = 0$ defines the field of parallel vector fields $v_i(x)$. The integrability conditions for these equations, that is, the existence of such parallel vectors was established in ref.[12]. The answer is that it should be possible to choose a coordinate system, in which the aforementioned conditions can be expressed as decomposability of the matrix G_{ij} , $1 \leq i, j \leq N$, into a direct sum of a constant diagonalizable matrix $G_{ij}^{(0)}$, $1 \leq i, j \leq s$, and a matrix $G_{ij}^{(1)}$, $s+1 \leq i, j \leq N$, depending only on the coordinates $x^{s+1}, \dots, x^N$, that is,

$$G = \begin{pmatrix} G_{11}^{(0)} & \dots & G_{1s}^{(0)} & 0 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ G_{s1}^{(0)} & \dots & G_{ss}^{(0)} & 0 & \dots & 0 \\ 0 & \dots & 0 & G_{s+1\ s+1}^{(1)} & \dots & G_{s+1\ N}^{(1)} \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \dots & 0 & G_{N\ s+1}^{(1)} & \dots & G_{N\ N}^{(1)} \end{pmatrix}. \quad (39)$$

The integer s , which is between 1 and N , is the number of linearly independent vectors $(v_i^{(1)}, \dots, v_i^{(s)})$ which generate a complete set of solutions of the system

$$\sum_n R_{nij}^l v^n = 0, \quad \sum_n R_{nij; m_1 \dots m_q}^l v^n = 0, \quad q \geq 0;$$

R_{nij}^l are the components of the Riemann tensor. (Note that the algebraic compatibility of these equations testifies to the existence of the parallel vector fields, for more details see ref.[13].) With these coordinates, the covariantly constant vectors $v_i^{(\ell)}$ are actually constant, and $v_i^{(\ell)} = 0$ for $s+1 \leq i \leq N$. Thus the covariantly constant vector $\Phi_{;j} \equiv \delta \Phi / \delta x^j$ satisfies

$$\delta \Phi / \delta x^i = \begin{cases} c_i, & \text{if } 1 \leq i \leq s \\ 0, & \text{if } s+1 \leq i \leq N, \end{cases} \quad (40)$$

where the c_i are constants. With those coordinates, $\Phi = \sum_1^s c_i \phi^i$ and Eq.37 becomes ($c^j = G^{(0)j i} c_i$)

$$\sum_{i=1}^s c^j \partial T / \partial x^j - T = 0. \quad (41)$$

The general solution of this equation is $T = \exp(\Phi / \sum_i c_i \phi^i) \tilde{T}$, where $\sum_i c^j \partial \tilde{T} / \partial x^j = 0$. One may verify that this equation is satisfied by the Toda theories[3], as well as by their non-abelian versions. In particular our black-hole solution is conformally invariant, at the classical level. However, it is easy to see that $e^{2\phi}$ transforms as a $(1, 1)$ operator while r and θ have vanishing weights. Thus the background metric Eq.10 is trivially invariant. More generally, Eq.40 shows that the dilaton field does not depend upon the variables

$x^s, \dots, x^N$ for which the background metric is non-trivial. Thus the latter is classically inert under conformal transformation. Clearly T plays the role of the 2D cosmological term.

Our next topic is the W algebraic structure of our black-hole solutions. As recalled at the beginning, the abelian Toda systems provide[3] Noether realizations of the W -algebras. It is interesting to note that for the non-abelian versions of the Toda systems, the W -algebras are basically different. In particular, for the case of B_2 , all three elements are of second order[16]. They may be written as (we consider one of the two holomorphic components, say, the z_+ one).

$$W_{\pm} = 2p_{\pm}p_0 - \partial_{\pm}p_{\pm}, \quad W_0 = p_0^2 + p_+p_- - \partial_+p_0 \quad (42)$$

where the fields $p_{\pm}, p_0$ are expressed as

$$p_{\pm} = e^{\pm\nu}(\pm \tanh(r)\partial_+\theta + \partial_+r), \quad p_0 = \partial_+\phi. \quad (43)$$

The field ν is defined by

$$\partial_+\nu = \frac{\cosh 2r}{\cosh^2 r}\partial_+\theta, \quad \partial_-\nu = \cosh^{-2}(r)\partial_-\theta \quad (44)$$

Note that it follows from the field equations that

$$\partial_-((\cosh 2r / \cosh^2 r)\partial_+\theta) = \partial_+(\cosh^{-2}(r)\partial_-\theta).$$

Thus Eqs.44 may be integrated. Using these relations, it is easy to get convinced that $\partial_-W_{\pm} = 0$, $\partial_-W_0 = 0$, so that they define three conserved quantities (characteristic integrals). One of them gives back the stress energy tensor as expected, since $\theta_{++} = 4W_0$, where $\theta_{\mu\nu}$ is given by Eq.2. In the same way as for the A_N abelian Toda case[18], we expect that the other conserved quantities will correspond to particular diffeomorphisms of the target space.

Another general point is that there exists[17] a supersymmetrized version of the dynamical system Eq.9. In this, Eq.21 is replaced by

$$\mathcal{D}_+ \left(\mathcal{D}_- \hat{g}_0^{-1} \hat{g}_0 \right) + \left[N_-, \hat{g}_0^{-1} N_+ \hat{g}_0 \right]_+ = 0. \quad (45)$$

Here one uses a classical Lie super-algebra supplied with a $\mathbf{Z}$ -gradation. $\mathcal{D}_{\pm}$ are covariant super-derivatives in $2|2$ -super-space; $\hat{g}_0$ is a function on the Grassmannian hull of the super-Lie-group generated by $\mathcal{G}_0$. $N_{\pm}$ are fixed (odd) elements of $\mathcal{G}_{\pm 1}$. This gives a supersymmetrized black hole.

In conclusion, this discussion was at the purely classical level, and the quantum case is left for further studies. We expect that the complete integrability will be of great help to quantize keeping conformal invariance, as it was already the case for the abelian Toda systems.

Acknowledgements

One of the authors (M. S.) is indebted to A. Leznov, Yu. Manin, A. Razumov, and P. Sorba for the useful discussions. He is also grateful to ENS LAPP in Lyon and LPTENS in Paris for kind hospitality. This work was partially supported by the European Community Twinning Program.

References

- [1] A.Belavin, A.Polyakov, A.Zamolodchikov: Nucl. Phys. **241B**, 333 (1984).
- [2] J.-L.Gervais, A.Neveu: Nucl. Phys. **B199** (1982) 59, **B209** (1982) 125, **B224** (1983) 329, **B238** (1984) 125, **B238** (1984) 396, **B257** [FS14] (1985) 59; Comm.Math. Phys. **100**, 15 (1985).
- [3] A.Bilal, J.-L.Gervais: Phys. Lett. **206B**,412 (1988); Nucl. Phys. **314B**,646 (1989), **318B**,579 (1989).
- [4] A.N.Leznov, M.V.Saveliev: Acta Appl. Math. **16**,1 (1989).
- [5] A.N.Leznov, M.V.Saveliev: Lett. Math. Phys. **6**,505 (1982); Comm. Math. Phys. **89**,59 (1983).
- [6] S. Elitsur, A. Forge, E. Rabinovici: *Some global aspects of string compactification*, preprint RI-141(90).
- [7] K. Bardakci, M. Crescimanno, E. Rabinovici: *Parafermions from Coset models*, LBL preprint (1990).
- [8] M. Rocek, K. Schoutens, A. Sevrin: *Off-shell WZNW models in extended superspace*, IASSNS-HEP-91/14.
- [9] G. Mandal, A. Sengupta, S. Wadia: *Classical solutions of 2-dimensional string theory* IASSYS-HEP-91/10.
- [10] E. Witten: Phys. Rev. **D44** (1991) 314, *On black holes in string theory*; Lecture at Strings '91, Stonybrook, June 1991.
- [11] P.Forgács, A.Wipf, J.Balog, L.Fehér, and L.O'Raifeartaigh: *Phys. Lett.* **227B** (1989) 214; J.Balog, L.Fehér, L.O'Raifeartaigh, P.Forgács and A.Wipf: *Ann. Phys.* (N.Y.) **203** (1990) 76; *Phys. Lett.* **244B** (1990) 435; L.O'Raifeartaigh and A.Wipf: *Phys. Lett.* **251B** (1990) 361; L.O'Raifeartaigh, P.Ruelle, I. Tsutsui and A.Wipf: *W-Algebras for Generalized Toda Theories*, Dublin preprint DIAS-STP-91-03, *Comm. Math. Phys.* to appear; L.Fehér, L.O'Raifeartaigh, P.Ruelle, I. Tsutsui and A.Wipf: *Generalized Toda theories and W-algebras associated with integer grading*, Dublin preprint DIAS-STP-91-17, *Ann. Phys.* to appear; L.O'Raifeartaigh, P.Ruelle and I. Tsutsui, *Phys. Lett.* **258B** (1990) 359.
- [12] L.P.Eisenhart: Thans. Amer. Math. Soc. **27**,563 (1925); P.A.Shirokov: Kazan Izv. Phys. Math. Soc. **(2) 25**,86 (1925) (in Russian).
- [13] L.P.Eisenhart:*Riemannian Geometry*, Princeton Univ. Press, Princeton, 1926.
- [14] A.N.Leznov, M.V.Saveliev: Comm. Math. Phys. **74** (1980) 111.
- [15] R. Gilmore: *Lie groups, Lie algebras, and some of their applications*, Wiley-Interscience Publication, 1974.

- [16] M.V. Saveliev: *Mod. Phys. Lett.* **5A** 27 (1990) 2223.
- [17] D.A. Leites, M.V. Saveliev, V.V. Serganova: *in Group Theoretical Methods in Physics* V.I. Manko, M.A. Markov, eds., VNU Sci. Press **1** (1986) 255.
- [18] J.-L. Gervais, Y. Matsuo: *Phys. Lett.* **274B** (1992) 309; A_n - W -geometry, LPTENS preprint 91/35.